



TG/UROCH(proj.10)

ORIGINAL: English

DATE: 2016-07-20

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS

Geneva

DRAFT

UROCHLOA *

UPOV Code: UROCH_RUZ; UROCH_DIC; UROCH_HUM;
UROCH_DEC; UROCH_BRI; UROCH_RBR; UROCH_RDB

Urochloa brizantha (Hochst. ex A. Rich.) R. D. Webster;
Urochloa decumbens (Stapf) R. D. Webster;
Urochloa dictyoneura (Fig. & De Not.) Veldkamp;
Urochloa humidicola (Rendle) Morrone & Zuloaga;
Urochloa ruziziensis (R. Germ. & C. M. Evrard) Crins;
Urochloa ruziziensis (R. Germ. & C. M. Evrard) Crins x *U. brizantha*
(Hochst. ex A. Rich.) R. D. Webster;
Urochloa ruziziensis x *Urochloa decumbens* x *Urochloa brizantha*

GUIDELINES

FOR THE CONDUCT OF TESTS

FOR DISTINCTNESS, UNIFORMITY AND STABILITY

prepared by (an) expert(s) from Brazil

to be considered by the

*Enlarged Editorial Committee at its meeting
to be held in Geneva, on January 11 and 12, 2017*

Disclaimer: this document does not represent UPOV policies or guidance

Alternative Names:*

Botanical name	English	French	German	Spanish
<i>Urochloa brizantha</i> (Hochst. ex A. Rich.) R. D. Webster, <i>Brachiaria brizantha</i> (Hochst. ex A. Rich.) Stapf, <i>Panicum brizanthum</i> Hochst. ex A. Rich.	Bread Grass, Palisade grass, Palisade signal grass, Signal Grass		Palisadengras	Pasto alambre, Pasto señal, Zacate señal, Zacate signal, Brachiaria

* These names were correct at the time of the introduction of these Test Guidelines but may be revised or updated. [Readers are advised to consult the UPOV Code, which can be found on the UPOV Website (www.upov.int), for the latest information.]

Alternative Names:*

<i>Botanical name</i>	<i>English</i>	<i>French</i>	<i>German</i>	<i>Spanish</i>
<i>Urochloa decumbens</i> (Stapf) R. D. Webster, <i>Brachiaria decumbens</i> Stapf	Basilisk signal grass, Signal grass, Spreading liverseed grass, Surinam grass		Surinamgras	Zacate Surinam, Pasto chontalpo, Pasto de la palizada, Pasto de las orillas, Pasto peludo, Pasto prodigio, Zacate prodigio, <i>Brachiaria</i>
<i>Urochloa dictyoneura</i> (Fig. & De Not.) Veldkamp , <i>Brachiaria dictyoneura</i> (Fig. & De Not.) Stapf, <i>Panicum dictyoneurum</i> Fig. & De Not.	Koronivia grass			
<i>Urochloa humidicola</i> (Rendle) Morrone & Zuloaga, <i>Brachiaria humidicola</i> (Rendle) Schweick.; <i>Panicum humidicola</i> Rendle	Creeping signal grass, Koronivia grass	Koronivia		Braquiaria dulce, Kikuyu de la Amazonía, Pasto humidicola, Pasto humidicola dulce
<i>Urochloa ruziziensis</i> (R. Germ. & C. M. Evrard) Crins, <i>Brachiaria ruziziensis</i> R. Germ. & C. M. Evrard	Congo grass, Congo signal grass, Ruzi grass			Congo señal, Gambutera, Kenia, Pasto Congo, Pasto ruzi, <i>Brachiaria</i>
<i>Urochloa ruziziensis</i> (R. Germ. & C. M. Evrard) Crins x <i>U. brizantha</i> (Hochst. ex A. Rich.) R. D. Webster, <i>Brachiaria ruziziensis</i> R. Germ. & C. M. Evrard x <i>B. brizantha</i> (Hochst. ex A. Rich.) Stapf				
<i>Urochloa ruziziensis</i> x <i>Urochloa decumbens</i> x <i>Urochloa brizantha</i> , <i>Brachiaria ruziziensis</i> x <i>Brachiaria decumbens</i> x <i>Brachiaria brizantha</i>				

The purpose of these guidelines ("Test Guidelines") is to elaborate the principles contained in the General Introduction (document TG/1/3), and its associated TGP documents, into detailed practical guidance for the harmonized examination of distinctness, uniformity and stability (DUS) and, in particular, to identify appropriate characteristics for the examination of DUS and production of harmonized variety descriptions.

ASSOCIATED DOCUMENTS

These Test Guidelines should be read in conjunction with the General Introduction and its associated TGP documents.

<u>TABLE OF CONTENTS</u>	<u>PAGE</u>
1. SUBJECT OF THESE TEST GUIDELINES.....	4
2. MATERIAL REQUIRED.....	4
3. METHOD OF EXAMINATION.....	4
3.1 NUMBER OF GROWING CYCLES	4
3.2 TESTING PLACE	4
3.3 CONDITIONS FOR CONDUCTING THE EXAMINATION.....	4
3.4 TEST DESIGN.....	4
3.5 ADDITIONAL TESTS.....	5
4. ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY	5
4.1 DISTINCTNESS	5
4.2 UNIFORMITY	6
4.3 STABILITY.....	6
5. GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL.....	6
6. INTRODUCTION TO THE TABLE OF CHARACTERISTICS	7
6.1 CATEGORIES OF CHARACTERISTICS	7
6.2 STATES OF EXPRESSION AND CORRESPONDING NOTES	7
6.3 TYPES OF EXPRESSION.....	7
6.4 EXAMPLE VARIETIES.....	7
6.5 LEGEND	8
7. TABLE OF CHARACTERISTICS/TABLEAU DES CARACTÈRES/MERKMALSTABELLE/TABLA DE CARACTERES	9
8. EXPLANATIONS ON THE TABLE OF CHARACTERISTICS.....	13
8.1 EXPLANATIONS COVERING SEVERAL CHARACTERISTICS	13
8.2 EXPLANATIONS FOR INDIVIDUAL CHARACTERISTICS	13
9. LITERATURE	17
10. TECHNICAL QUESTIONNAIRE.....	18

1. Subject of these Test Guidelines

These Test Guidelines apply to all varieties of *Urochloa brizantha* (Hochst. ex A. Rich.) R. D. Webster, *Urochloa decumbens* (Stapf) R. D. Webster, *Urochloa dictyoneura* (Fig. & De Not.) Veldkamp, *Urochloa humidicola* (Rendle) Morrone & Zuloaga, *Urochloa ruziziensis* (R. Germ. & C. M. Evrard) Crins, *Urochloa ruziziensis* (R. Germ. & C. M. Evrard) Crins x *U. brizantha* (Hochst. ex A. Rich.) R. D. Webster, *Urochloa ruziziensis* x *Urochloa decumbens* x *Urochloa brizantha*.

2. Material Required

2.1 The competent authorities decide on the quantity and quality of the plant material required for testing the variety and when and where it is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must ensure that all customs formalities and phytosanitary requirements are complied with.

2.2 The material is to be supplied in the form of seed.

2.3 The minimum quantity of plant material, to be supplied by the applicant, should be:

500 g of seed.

The seed should meet the minimum requirements for germination, species and analytical purity, health and moisture content, specified by the competent authority. In cases where the seed is to be stored, the germination capacity should be as high as possible and should, be stated by the applicant.

2.4 The plant material supplied should be visibly healthy, not lacking in vigor, nor affected by any important pest or disease.

2.5 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

3. Method of Examination

3.1 *Number of Growing Cycles*

3.1.1 The minimum duration of tests should normally be two independent growing cycles.

3.1.2 The two independent growing cycles should be in the form of two separate plantings.

3.2 *Testing Place*

Tests are normally conducted at one place. In the case of tests conducted at more than one place, guidance is provided in TGP/9 "Examining Distinctness".

3.3 *Conditions for Conducting the Examination*

The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.

3.4 *Test Design*

3.4.1 For apomictic varieties, each test should be designed to result in a total of at least 40 spaced plants which should be divided between at least 2 replicates.

3.4.2 For cross-pollinated varieties, each test should be designed to result in a total of at least 60 spaced plants which should be divided between at least 3 replicates.

3.4.3 The design of the tests should be such that plants or parts of plants may be removed for measurement or counting without prejudice to the observations which must be made up to the end of the growing cycle.

3.5 *Additional Tests*

Additional tests, for examining relevant characteristics, may be established.

4. Assessment of Distinctness, Uniformity and Stability

4.1 *Distinctness*

4.1.1 General Recommendations

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

4.1.2 Consistent Differences

The differences observed between varieties may be so clear that more than one growing cycle is not necessary. In addition, in some circumstances, the influence of the environment is not such that more than a single growing cycle is required to provide assurance that the differences observed between varieties are sufficiently consistent. One means of ensuring that a difference in a characteristic, observed in a growing trial, is sufficiently consistent is to examine the characteristic in at least two independent growing cycles.

4.1.3 Clear Differences

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Test Guidelines are familiar with the recommendations contained in the General Introduction prior to making decisions regarding distinctness.

4.1.4 Number of Plants / Parts of Plants to be Examined

In the case of apomictic varieties, unless otherwise indicated, for the purposes of distinctness, all observations on single plants should be made on 20 plants or parts taken from each of 20 plants and any other observations made on all plants in the test, disregarding any off-type plants.

In the case of cross-pollinated varieties, unless otherwise indicated, for the purposes of distinctness, all observations on single plants should be made on 60 plants or parts taken from each of 60 plants and any other observations made on all plants in the test, disregarding any off-type plants.

4.1.5 Method of Observation

The recommended method of observing the characteristic for the purposes of distinctness is indicated by the following key in the second column of the Table of Characteristics (see document TGP/9 "Examining Distinctness", Section 4 "Observation of characteristics"):

MG: single measurement of a group of plants or parts of plants
MS: measurement of a number of individual plants or parts of plants
VG: visual assessment by a single observation of a group of plants or parts of plants
VS: visual assessment by observation of individual plants or parts of plants

Type of observation: visual (V) or measurement (M)

"Visual" observation (V) is an observation made on the basis of the expert's judgment. For the purposes of this document, "visual" observation refers to the sensory observations of the experts and, therefore, also includes smell, taste and touch. Visual observation includes observations where the expert uses reference points (e.g. diagrams, example varieties, side-by-side

comparison) or non-linear charts (e.g. color charts). Measurement (M) is an objective observation against a calibrated, linear scale e.g. using a ruler, weighing scales, colorimeter, dates, counts, etc.

Type of record: for a group of plants (G) or for single, individual plants (S)

For the purposes of distinctness, observations may be recorded as a single record for a group of plants or parts of plants (G), or may be recorded as records for a number of single, individual plants or parts of plants (S). In most cases, “G” provides a single record per variety and it is not possible or necessary to apply statistical methods in a plant-by-plant analysis for the assessment of distinctness.

In cases where more than one method of observing the characteristic is indicated in the Table of Characteristics (e.g. VG/MG), guidance on selecting an appropriate method is provided in document TGP/9, Section 4.2.

4.2 *Uniformity*

4.2.1 It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding uniformity. However, the following points are provided for elaboration or emphasis in these Test Guidelines:

4.2.2 The assessment of uniformity for cross-pollinated varieties should be according to the recommendations for cross-pollinated varieties in the General Introduction.

4.2.3 For the assessment of uniformity of apomictic varieties, a population standard of 2% and an acceptance probability of at least 95 % should be applied. In the case of a sample size of 40 plants, 2 off-types are allowed.

4.3 *Stability*

4.3.1 In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.

4.3.2 Where appropriate, or in cases of doubt, stability may be further examined by testing a new seed stock to ensure that it exhibits the same characteristics as those shown by the initial material supplied.

5. Grouping of Varieties and Organization of the Growing Trial

5.1 The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness are aided by the use of grouping characteristics.

5.2 Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.

5.3 The following have been agreed as useful grouping characteristics:

- (a) Leaf blade: hairiness (characteristic 11)
- (b) Inflorescence: shape of rachis in cross section (characteristic 17)
- (c) Flower: stigma color (characteristic 19)

5.4 Guidance for the use of grouping characteristics, in the process of examining distinctness, is provided through the General Introduction and document TGP/9 “Examining Distinctness”.

6. Introduction to the Table of Characteristics

6.1 *Categories of Characteristics*

6.1.1 Standard Test Guidelines Characteristics

Standard Test Guidelines characteristics are those which are approved by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular circumstances.

6.1.2 Asterisked Characteristics

Asterisked characteristics (denoted by *) are those included in the Test Guidelines which are important for the international harmonization of variety descriptions and should always be examined for DUS and included in the variety description by all members of the Union, except when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.

6.2 *States of Expression and Corresponding Notes*

6.2.1 States of expression are given for each characteristic to define the characteristic and to harmonize descriptions. Each state of expression is allocated a corresponding numerical note for ease of recording of data and for the production and exchange of the description.

6.2.2 In the case of qualitative and pseudo-qualitative characteristics (see Chapter 6.3), all relevant states of expression are presented in the characteristic. However, in the case of quantitative characteristics with 5 or more states, an abbreviated scale may be used to minimize the size of the Table of Characteristics. For example, in the case of a quantitative characteristic with 9 states, the presentation of states of expression in the Test Guidelines may be abbreviated as follows:

State	Note
small	3
medium	5
large	7

However, it should be noted that all of the following 9 states of expression exist to describe varieties and should be used as appropriate:

State	Note
very small	1
very small to small	2
small	3
small to medium	4
medium	5
medium to large	6
large	7
large to very large	8
very large	9

6.2.3 Further explanation of the presentation of states of expression and notes is provided in document TGP/7 "Development of Test Guidelines".

6.3 *Types of Expression*

An explanation of the types of expression of characteristics (qualitative, quantitative and pseudo-qualitative) is provided in the General Introduction.

6.4 *Example Varieties*

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic.

6.5 *Legend*

(*) Asterisked characteristic – see Chapter 6.1.2

QL Qualitative characteristic – see Chapter 6.3

QN Quantitative characteristic – see Chapter 6.3

PQ Pseudo-qualitative characteristic – see Chapter 6.3

MG, MS, VG, VS – see Chapter 4.1.5

(a)-(b) See Explanations on the Table of Characteristics in Chapter 8.

(+) See Explanations on the Table of Characteristics in Chapter 8.

7. Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteres

		English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
1. (*) (+)	VG	Plant: growth habit	Plante : port	Pflanze: Wuchsform	Planta: porte		
QN		erect	dressé	aufrecht	erecto	BRS Piatã, CIAT BR02/1718	1
		semi erect	demi-dressé	halbaufrecht	semierecto	BRS Tupi, Llanero	3
		semi prostate	demi-étalé	halbliegend	semipostrado	MIXE LN 45, Mulato II	5
		prostrate	étalé	liegend	postrado	Humidicola comum	7
2. (*) (+)	MS	Plant: height	Plante : hauteur	Pflanze: Höhe	Planta: altura		
QN	(a)	short	basse	niedrig	corta	BRS Tupi	3
		medium	moyenne	mittel	media	BRS Piatã, MIXE LN 45, Mulato II	5
		tall	haute	hoch	alta	CIAT BR02/1718, Xaraés	7
3. (*) (+)	MS	Stolon: length of internode	Stolon : longueur de l'entre-nœud	Ausläufer: Internodienlänge	Estolón: longitud del entrenudo		
QN		absent or very short	absent ou très court	fehlend oder sehr kurz	ausente o muy corto	BRS Piatã	1
		short	court	kurz	corto	Mulato II	3
		medium	moyen	mittel	medio	Humidicola comum	5
		long	long	lang	largo	BRS Tupi	7
4. (*) (+)	MS	Culm: length of internode	Tige : longueur de l'entre-nœud	Halm: Internodienlänge	Macollo: longitud del entrenudo		
QN		short	court	kurz	corto	BRS Tupi	3
		medium	moyen	mittel	medio	MIXE LN 45	5
		long	long	lang	largo	Xaraés	7
5. (+)	MS	Culm: diameter	Tige : diamètre	Halm: Durchmesser	Macollo: diámetro		
QN		small	petit	klein	pequeño		1
		medium	moyen	mittel	medio	MIXE LN 45, Mulato II	2
		large	grand	groß	grande		3
6. (*) (+)	VG	Flag leaf: curvature	Dernière feuille : courbure	Fahnenblatt: Biegung	Última hoja: curvatura		
QN	(b)	absent or weak	absente ou faible	fehlend oder gering	ausente o débil		1
		medium	moyenne	mittel	media		2
		strong	forte	stark	fuerte		3
7. (*) (+)	VG	Flag leaf: width	Dernière feuille : largeur	Fahnenblatt: Breite	Última hoja: anchura		
QN	(b)	narrow	étroite	schmal	estrecha	BRS Piatã	1
		medium	moyenne	mittel	media	MIXE LN 45, Mulato II	2
		broad	large	breit	ancha		3

		English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
8. (*)	VG	Flag leaf: distribution of hairs on sheath	Dernière feuille : répartition des poils sur la gaine	Fahnenblatt: Verteilung der Haare auf Blattscheide	Última hoja: distribución de los pelos en la vaina		
PQ	(b)	at base	à la base	an der Basis	en la base		1
		at apex	au sommet	an der Spitze	en el ápice	MIXE LN 45	2
		on margins	en bordure	an den Rändern	en los márgenes		3
		throughout	partout	überall	en la totalidad	BRS Piatã, Mulato II	4
9.	MS	Leaf blade: length	Limbe : longueur	Blattspreite: Länge	Limbo: longitud		
QN	(b)	short	court	kurz	corto	Basilisk, Humidícola comum	3
		medium	moyen	mittel	medio	MIXE LN 45	5
		long	long	lang	largo	BRS Piatã, Mulato II	7
10. (*)	MS	Leaf blade: width	Limbe : largeur	Blattspreite: Breite	Limbo: anchura		
QN	(b)	narrow	étroit	schmal	estrechas	BRS Piatã	3
		medium	moyen	mittel	medio	MIXE LN 45	5
		broad	large	breit	anchas	Mulato II	7
11. (*)	VG	Leaf blade: hairiness	Limbe : pilosité	Blattspreite: Behaarung	Limbo: vellosoidad		
QL	(b)	absent	absente	fehlend	ausente	BRS Tupi	1
		present	présente	vorhanden	presente	Mulato II	9
12. (*)	VG	Leaf blade: distribution of hairs	Limbe : répartition des poils	Blattspreite: Verteilung der Haare	Limbo: distribución de los pelos		
PQ	(b)	on upper surface only	uniquement sur la face supérieure	nur auf Oberseite	solo en el haz	BRS Tupi, Llanero	1
		on lower surface only	uniquement sur la face inférieure	nur auf Unterseite	solo en el envés	MIXE LN 45	2
		on margins only	uniquement en bordure	nur auf Rändern	solo en los márgenes	Marandú, Xaraés	3
		on both surfaces	sur les deux faces	auf beiden Seiten	en ambas superficies	Mulato II, Basilisk	4
13. (*)	VG	Leaf sheath: density of hairs	Gaine de la feuille: densité de la pilosité	Blattscheide: Dichte der Behaarung	Vaina de la hoja: densidad de la vellosoidad		
QN	(b)	absent or sparse	absente ou épars	fehlend oder locker	ausente o escasa	BRS Piatã	1
		medium	moyenne	mittel	media		2
		dense	dense	dicht	densa	Mulato II	3
14. (*) (+)	MS	Inflorescence: length of peduncle	Inflorescence : longueur du pédoncule	Blütenstand: Länge des Blütenstandsstiels	Inflorescencia: longitud del pedúnculo		
QN	(a)	short	court	kurz	corto		3
		medium	moyen	mittel	medio	Mulato II	5
		long	long	lang	largo	BRS Piatã, MIXE LN 45	7

		English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
15.	MS	Inflorescence: length of rachis	Inflorescence : longueur du rachis	Blütenstand: Länge der Spindeln	Inflorescencia: longitud del raquis		
(+)							
QN	(a)	short	court	kurz	corto	Mulato II	3
		medium	moyen	mittel	medio	Llanero	5
		long	long	lang	largo	Marandú	7
16.	MS	Inflorescence: length of basal racemes	Inflorescence : longueur des racèmes de la base	Blütenstand: Länge der basalen Blütentrauben	Inflorescencia: longitud de los racimos basales		
(+)							
QN	(a)	short	courts	kurz	cortos	BRS Tupi, Humidícola comum	3
		medium	moyens	mittel	medios	MIXE LN 45, Mulato II	5
		long	longs	lang	largos	Marandú	7
17.	VG	Inflorescence: shape of rachis in cross section	Inflorescence : forme du rachis en section transversale	Blütenstand: Form der Spindeln im Querschnitt	Inflorescencia: forma del raquis en sección transversal		
(*)							
(+)							
PQ		triangular	triangulaire	dreieckig	triangular	MIXE LN 45	1
		winged	ailée	geflügelt	alado	Mulato II	2
		crescent	en croissant	halbmondförmig	lunular	BRS Piatã	3
18.	VG	Spikelet: pubescence	Épillet : pubescence	Ährchen: Behaarung	Espiguilla: pubescencia		
(*)							
QN	(a)	absent or very sparse	absente ou très épars	fehlend oder sehr locker	ausente o muy escasa	BRS Piatã	1
		sparse	épars	locker	escasa	Humidícola comum	3
		medium	moyenne	mittel	media	Mulato II, Xaraés	5
		dense	dense	dicht	densa	BRS Tupi, Llanero	7
19.	VG	Flower: stigma color	Fleur : couleur des stigmates	Blüte: Farbe der Narbe	Flor: color del estigma		
(*)							
(+)							
PQ		white	blanc	weiß	blanco	Mulato II	1
		light purple	violet clair	hellpurpurn	púrpura claro	Llanero	2
		medium purple	violet moyen	mittelpurpurn	púrpura medio	BRS Piatã, MIXE LN 45	3
		dark purple	violet foncé	dunkelpurpurn	púrpura oscuro	Marandú, Toledo	4
20.	VG	Glume: anthocyanin coloration	Glume : pigmentation anthocyanique	Hüllspelze: Anthocyanfärbung	Gluma: pigmentación antocianica		
(+)							
QN		absent or very weak	absente ou très faible	fehlend oder sehr gering	ausente o muy débil	BRS Piatã	1
		weak	absente	gering	débil	Basilisk	3
		medium	moyenne	mittel	media	Marandú	5
		strong	forte	stark	fuerte	Llanero	7

		English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
21. (*) (+)	MG	Time of beginning of flowering	Époque du début de floraison	Zeitpunkt des Blühbeginns	Época de inicio de la floración		
	QN	early	précoce	früh	temprana	BRS Piatã, Basilisk, Llanero	3
		medium	moyenne	mittel	media	Marandú	5
		late	tardive	spät	tardía	Xaraés	7

8. Explanations on the Table of Characteristics


8.1 *Explanations covering several characteristics*

Characteristics containing the following key in the second column of the Table of Characteristics should be examined as indicated below:

- (a) Observations should be made at the time of beginning of flowering.
- (b) Observations on fully developed leaves should be made on the penultimate leaf of the main culm.

8.2 *Explanations for individual characteristics*

Ad. 1: Plant: growth habit



Ad. 2: Plant: height


The height of the plant should be measured in the center of the plant, from the first leaf below the flag leaf to the ground level, excluding the inflorescence.




Ad. 4: Culm: length of internode

Ad. 5: Culm: diameter


The assessment of the length of internode should be made in the middle third of plant; it does not refer to floral culm.




Ad. 6: Flag leaf: curvature



1
absent or weak




2
medium



3
strong


Ad. 7: Flag leaf: width




Ad. 14: Inflorescence: length of peduncle

Ad. 15: Inflorescence: length of rachis

Ad. 16: Inflorescence: length of basal racemes



Ad. 17: Inflorescence: shape of rachis in cross section



Ad. 19: Flower: stigma color

To be observed at anthesis.

Ad. 20: Glume: anthocyanin coloration

The anthocyanin coloration on the glume should be assessed at the beginning of flowering, when 50% of the plants have at least one inflorescence fully emerged and before the opening of flowers.

Ad. 21: Time of beginning of flowering

The time of beginning of flowering should be assessed when 50% of the plants have at least one inflorescence fully emerged.

9. Literature

Assis, G.M.L. de, Euclides, R.F., Cruz, C.D. and Valle, C. B. do. 2003: Discriminação de Espécies de Brachiaria Baseada em Diferentes Grupos de Caracteres Morfológicos. R. Bras. Zootec., v.32, n.3, pp.576-584

Dahmer, N., Schifino-Wittman, M.T., Dall'Agnol, M., Castro, B de, 2008: Cytogenetic data for Paspalum notatum Flüge accessions. Sci. Agric., Piracicaba, v.65, n.4, p.381-388.

Miles, J. W., Maass, B. L. and Valle, C. B. do. eds., 1996: Brachiaria: Biology, Agronomy, and Improvement. CIAT Publication No. 259

Pozzobon, M.T., Valls, J.M., 1997: Chromosome number in germplasm accessions of Paspalum notatum (Gramineae). Braz. J. Genet., Ribeirão preto, v.20, n.1, p.29-34

Simioni, C., Schifino-Wittman, M.T., Dall'Agnol, M., 2006: Sexual polyploidization in red clover, Sci. Agric., Piracicaba, v.63, n.1, p.26-31

10. Technical Questionnaire

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

	Application date:
	(not to be filled in by the applicant)

TECHNICAL QUESTIONNAIRE to be completed in connection with an application for plant breeders' rights		
1. Subject of the Technical Questionnaire		
1.1.1 Botanical name	<i>Urochloa decumbens</i> (Stapf) R. D. Webster	[]
1.1.2 Common name	Basilisk signal grass, Signal grass, Spreading liverseed grass, Surinam grass	
1.2.1 Botanical name	<i>Urochloa humidicola</i> (Rendle) Morrone & Zuloaga	[]
1.2.2 Common name	Creeping signal grass, Koronivia grass	
1.3.1 Botanical name	<i>Urochloa ruziziensis</i> (R. Germ. & C. M. Evrard) Morrone & Zuloaga	[]
1.3.2 Common name	Congo grass, Congo signal grass, Ruzi grass	
1.4.1 Botanical name	<i>Urochloa dictyoneura</i> (Fig. & De Not.) Veldkamp	[]
1.4.2 Common name	Koronivia grass	
1.5.1 Botanical name	<i>Urochloa brizantha</i> (Hochst. ex A. Rich.) R. D. Webster	[]
1.5.2 Common name	Bread Grass, Palisade grass, Palisade signal grass, Signal Grass	
1.6.1 Botanical name	<i>Urochloa ruziziensis</i> (R. Germ. & C. M. Evrard) Crins x <i>U. brizantha</i> (Hochst. ex A. Rich.) R. D. Webster	[]
1.6.2 Common name		
1.7.1 Botanical name	<i>Urochloa ruziziensis</i> x <i>Urochloa decumbens</i> x <i>Urochloa brizantha</i>	[]
1.7.2 Common name		

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

2. Applicant

Name

Address

Telephone No.

Fax No.

E-mail address

Breeder (if different from applicant)

3. Proposed denomination and breeder's reference

Proposed denomination
(if available)

Breeder's reference

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

#4. Information on the breeding scheme and propagation of the variety

4.1 Breeding scheme

Variety resulting from:

4.1.1 Crossing

(a) controlled cross []
(please state parent varieties)

(.....) x (.....)
female parent male parent

(b) partially known cross []
(please state known parent variety(ies))

(.....) x (.....)
female parent male parent

(c) unknown cross []

4.1.2 Mutation []
(please state parent variety)

4.1.3 Discovery and development []
(please state where and when discovered and how developed)

4.1.4 Other []
(please provide details)

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

4.2 Method of propagating the variety

4.2.1 Seed-propagated varieties

- | | | |
|-----|--------------------------|-----|
| (a) | apomictic | [] |
| (b) | non-apomictic | [] |
| (c) | Other | [] |
| | (please provide details) | |

--

4.2.2	Other	
	(please provide details)	[...]

--

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the note which best corresponds).

Characteristics	Example Varieties	Note
5.1 (11) Leaf blade: hairiness		
absent	BRS Tupi	1 []
present	Mulato II	9 []
5.2 (17) Inflorescence: shape of rachis in cross section		
triangular	MIXE LN 45	1 []
winged	Mulato II	2 []
crescent	BRS Piatã	3 []
5.3 (19) Flower: stigma color		
white	Mulato II	1 []
light purple	Llanero	2 []
medium purple	BRS Piatã, MIXE LN 45	3 []
dark purple	Marandú, Toledo	4 []

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

6. Similar varieties and differences from these varieties

Please use the following table and box for comments to provide information on how your candidate variety differs from the variety (or varieties) which, to the best of your knowledge, is (or are) most similar. This information may help the examination authority to conduct its examination of distinctness in a more efficient way.

Denomination(s) of variety(ies) similar to your candidate variety	Characteristic(s) in which your candidate variety differs from the similar variety(ies)	Describe the expression of the characteristic(s) for the similar variety(ies)	Describe the expression of the characteristic(s) for your candidate variety
<i>Example</i>	<i>Flower: stigma color</i>	<i>dark purple</i>	<i>light purple</i>

Comments:

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
#7. Additional information which may help in the examination of the variety 7.1 In addition to the information provided in sections 5 and 6, are there any additional characteristics which may help to distinguish the variety? Yes [] No [] (If yes, please provide details) 7.2 Are there any special conditions for growing the variety or conducting the examination? Yes [] No [] (If yes, please provide details) 7.3 Other information Ploidy []		
8. Authorization for release (a) Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health? Yes [] No [] (b) Has such authorization been obtained? Yes [] No [] If the answer to (b) is yes, please attach a copy of the authorization.		

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

9. Information on plant material to be examined or submitted for examination

9.1 The expression of a characteristic or several characteristics of a variety may be affected by factors, such as pests and disease, chemical treatment (e.g. growth retardants or pesticides), effects of tissue culture, different rootstocks, scions taken from different growth phases of a tree, etc.

9.2 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If the plant material has undergone such treatment, full details of the treatment must be given. In this respect, please indicate below, to the best of your knowledge, if the plant material to be examined has been subjected to:

- | | | | |
|-----|---|---------|--------|
| (a) | Microorganisms (e.g. virus, bacteria, phytoplasma) | Yes [] | No [] |
| (b) | Chemical treatment (e.g. growth retardant, pesticide) | Yes [] | No [] |
| (c) | Tissue culture | Yes [] | No [] |
| (d) | Other factors | Yes [] | No [] |

Please provide details for where you have indicated "yes".

.....

10. I hereby declare that, to the best of my knowledge, the information provided in this form is correct:

Applicant's name

Signature

Date

[End of document]