



TG/197/2(proj.3)

ORIGINAL: English

DATE: 2021-04-23

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS

Geneva

DRAFT

EUSTOMA

UPOV Code(s): EUSTO_GRA

Eustoma exaltatum (L.) Salisb. ex G. Don
subsp. *russellianum* (Hook.) Kartesz

GUIDELINES

FOR THE CONDUCT OF TESTS

FOR DISTINCTNESS, UNIFORMITY AND STABILITY

*prepared by experts from Japan
to be considered by the
Technical Working Party for Ornamental Plants and Forest Trees
at its fifty-third session, to be held in Roelofarendsveen, Netherlands,
from 2021-06-07 to 2021-06-11*

Disclaimer: this document does not represent UPOV policies or guidance

Alternative names:*

Botanical name	English	French	German	Spanish
<i>Eustoma exaltatum</i> (L.) Salisb. ex G. Don subsp. <i>russellianum</i> (Hook.) Kartesz, <i>Bilamista grandiflora</i> Raf., <i>Eustoma</i> <i>grandiflorum</i> (Raf.) Shinners, <i>Eustoma</i> <i>russellianum</i> (Hook.) G. Don, <i>Lisianthus</i> <i>russellianus</i> Hook.	Eustoma, Lisianthus	Eustoma, Lisianthus	Eustoma, Lisianthus	Eustoma, Lisianthus

The purpose of these guidelines ("Test Guidelines") is to elaborate the principles contained in the General Introduction (document TG/1/3), and its associated TGP documents, into detailed practical guidance for the harmonized examination of distinctness, uniformity and stability (DUS) and, in particular, to identify appropriate characteristics for the examination of DUS and production of harmonized variety descriptions.

ASSOCIATED DOCUMENTS

These Test Guidelines should be read in conjunction with the General Introduction and its associated TGP documents.

TABLE OF CONTENTS	PAGE
1. SUBJECT OF THESE TEST GUIDELINES.....	4
2. MATERIAL REQUIRED.....	4
3. METHOD OF EXAMINATION.....	4
3.1 Number of Growing Cycles.....	4
3.2 Testing Place.....	4
3.3 Conditions for Conducting the Examination.....	4
3.4 Test Design.....	6
3.5 Additional Tests.....	6
4. ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY.....	6
4.1 Distinctness.....	6
4.2 Uniformity.....	7
4.3 Stability.....	7
5. GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL.....	9
6. INTRODUCTION TO THE TABLE OF CHARACTERISTICS.....	10
6.1 Categories of Characteristics.....	10
6.2 States of Expression and Corresponding Notes.....	10
6.3 Types of Expression.....	10
6.4 Example Varieties.....	10
6.5 Legend.....	11
7. TABLE OF CHARACTERISTICS/TABLEAU DES CARACTÈRES/MERKMALSTABELLE/TABLA DE CARACTERES.....	12
8. EXPLANATIONS ON THE TABLE OF CHARACTERISTICS.....	20
8.1 Explanations covering several characteristics.....	20
8.2 Explanations for individual characteristics.....	21
9. LITERATURE.....	30
10 TECHNICAL QUESTIONNAIRE.....	31

1. Subject of these Test Guidelines

- 1.1 These Test Guidelines apply to all varieties of *Eustoma exaltatum* (L.) Salisb. ex G. Don subsp. *russellianum* (Hook.) Kartesz and interspecific hybrids.
- 1.2 Guidance on the use of Test Guidelines for interspecific hybrids that are not explicitly covered by Test Guidelines is provided in document TGP/13 "Guidance for New Types and Species."

2. Material Required

- 2.1 The competent authorities decide on the quantity and quality of the plant material required for testing the variety and when and where it is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must ensure that all customs formalities and phytosanitary requirements are complied with.
- 2.2 The material is to be supplied in the form of plants or seed.
- 2.3 The minimum quantity of plant material, to be supplied by the applicant, should be:

vegetatively propagated varieties: 20 plants
seed-propagated varieties: a sufficient quantity of seed to produce 40 plants.

In the case of seed, the seed should meet the minimum requirements for germination, species and analytical purity, health and moisture content, specified by the competent authority. In cases where the seed is to be stored, the germination capacity should be as high as possible and should, be stated by the applicant.

- 2.4 The plant material supplied should be visibly healthy, not lacking in vigor, nor affected by any important pest or disease.
- 2.5 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

3. Method of Examination

3.1 *Number of Growing Cycles*

- 3.1.1 The minimum duration of tests should normally be a single growing cycle.
- 3.1.2 The testing of a variety may be concluded when the competent authority can determine with certainty the outcome of the test.

3.2 *Testing Place*

Tests are normally conducted at one place. In the case of tests conducted at more than one place, guidance is provided in TGP/9 "Examining Distinctness".

3.3 *Conditions for Conducting the Examination*

- 3.3.1 The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.
- 3.3.2 Because daylight varies, color determinations made against a color chart should be made either in a suitable cabinet providing artificial daylight or in the middle of the day in a room without direct sunlight. The spectral distribution of the illuminant for artificial daylight should conform with the CIE Standard of Preferred Daylight D 6500 and should fall within the tolerances set out in the British Standard 950, Part I. These determinations should be made with the plant part placed against a white background. The color chart and version used should be specified in the variety description.

3.4 *Test Design*

- 3.4.1 In the case of vegetatively propagated varieties, each test should be designed to result in a total of at least 20 plants.
- 3.4.2 In the case of seed-propagated varieties, each test should be designed to result in a total of at least 40 plants.
- 3.4.3 The design of the tests should be such that plants or parts of plants may be removed for measurement or counting without prejudice to the observations which must be made up to the end of the growing cycle.

3.5 *Additional Tests*

Additional tests, for examining relevant characteristics, may be established.

4. Assessment of Distinctness, Uniformity and Stability

4.1 *Distinctness*

4.1.1 General Recommendations

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

4.1.2 Consistent Differences

The differences observed between varieties may be so clear that more than one growing cycle is not necessary. In addition, in some circumstances, the influence of the environment is not such that more than a single growing cycle is required to provide assurance that the differences observed between varieties are sufficiently consistent. One means of ensuring that a difference in a characteristic, observed in a growing trial, is sufficiently consistent is to examine the characteristic in at least two independent growing cycles.

4.1.3 Clear Differences

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Test Guidelines are familiar with the recommendations contained in the General Introduction prior to making decisions regarding distinctness.

4.1.4 Number of Plants or Parts of Plants to be Examined

In the case of vegetatively propagated varieties, unless otherwise indicated, for the purposes of distinctness, all observations on single plants should be made on 10 plants or parts taken from each of 10 plants and any other observation made on all plants in the test, disregarding any off-type plants.

In the case of seed-propagated varieties, unless otherwise indicated, for the purposes of distinctness, all observations on single plants should be made on 20 plants or parts taken from each of 20 plants and any other observation made on all plants in the test, disregarding any off-type plants.

4.1.5 Method of Observation

The recommended method of observing the characteristic for the purposes of distinctness is indicated by the following key in the Table of Characteristics (see document TGP/9 "Examining Distinctness", Section 4 "Observation of characteristics"):

MG: single measurement of a group of plants or parts of plants

MS: measurement of a number of individual plants or parts of plants

VG: visual assessment by a single observation of a group of plants or parts of plants

VS: visual assessment by observation of individual plants or parts of plants

Type of observation: visual (V) or measurement (M)

"Visual" observation (V) is an observation made on the basis of the expert's judgment. For the purposes of this document, "visual" observation refers to the sensory observations of the experts and, therefore, also includes smell, taste and touch. Visual observation includes observations where the expert uses reference points (e.g. diagrams, example varieties, side-by-side comparison) or non-linear charts (e.g. color charts). Measurement (M) is an objective observation against a calibrated, linear scale e.g. using a ruler, weighing scales, colorimeter, dates, counts, etc.

Type of record: for a group of plants (G) or for single, individual plants (S)

For the purposes of distinctness, observations may be recorded as a single record for a group of plants or parts of plants (G), or may be recorded as records for a number of single, individual plants or parts of plants (S). In most cases, "G" provides a single record per variety and it is not possible or necessary to apply statistical methods in a plant-by-plant analysis for the assessment of distinctness.

In cases where more than one method of observing the characteristic is indicated in the Table of Characteristics (e.g. VG/MG), guidance on selecting an appropriate method is provided in document TGP/9, Section 4.2.

4.2 *Uniformity*

4.2.1 It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding uniformity. However, the following points are provided for elaboration or emphasis in these Test Guidelines:

4.2.2 These Test Guidelines have been developed for the examination of vegetatively propagated and seed-propagated varieties. For varieties with other types of propagation, the recommendations in the General Introduction and document TGP/13 "Guidance for new types and species" Section 4.5 "Testing Uniformity" should be followed.

4.2.3 The assessment of uniformity for cross-pollinated varieties should be according to the recommendations for cross-pollinated varieties in the General Introduction.

4.2.4 The assessment of uniformity for hybrid varieties depends on the type of hybrid and should be according to the recommendations for hybrid varieties in the General Introduction.

4.2.5 For the assessment of uniformity of vegetatively propagated varieties and self-pollinated varieties, a population standard of 2% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 20 plants, 2 off-types are allowed.

4.3 *Stability*

4.3.1 In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.

4.3.2 Where appropriate, or in cases of doubt, stability may be further examined by testing a new seed or plant stock to ensure that it exhibits the same characteristics as those shown by the initial material supplied.

5. Grouping of Varieties and Organization of the Growing Trial
- 5.1 The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness are aided by the use of grouping characteristics.
- 5.2 Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.
- 5.3 The following have been agreed as useful grouping characteristics:
- (a) Plant: height (characteristic 1)
 - (b) Flower: type (characteristic 14)
 - (c) Flower: width (characteristic 17)
 - (d) Petal: main color of inner side (exclud part of base) (characteristic 27)
with the following groups
 - Gr. 1: white
 - Gr. 2: light green
 - Gr. 3: yellow
 - Gr. 4: orange
 - Gr. 5: pink
 - Gr. 6: red
 - Gr. 7: purple
 - Gr. 8: blue purple
 - (e) Petal: secondary color of inner side (characteristic 28)
with the following groups
 - Gr. 1: white
 - Gr. 2: light green
 - Gr. 3: yellow
 - Gr. 4: orange
 - Gr. 5: pink
 - Gr. 6: red
 - Gr. 7: purple
 - Gr. 8: blue purple
 - (f) Petal: distribution of secondary color of inner side (characteristic 29)
 - (g) Petal: color of base of inner side (characteristic 31)
 - (h) Only seed -propagated varieties: Time of beginning of flowering (characteristic 34)
- 5.4 Guidance for the use of grouping characteristics, in the process of examining distinctness, is provided through the General Introduction and document TGP/9 "Examining Distinctness".

6. Introduction to the Table of Characteristics

6.1 *Categories of Characteristics*

6.1.1 Standard Test Guidelines Characteristics

Standard Test Guidelines characteristics are those which are approved by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular circumstances.

6.1.2 Asterisked Characteristics

Asterisked characteristics (denoted by *) are those included in the Test Guidelines which are important for the international harmonization of variety descriptions and should always be examined for DUS and included in the variety description by all members of the Union, except when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.

6.2 *States of Expression and Corresponding Notes*

6.2.1 States of expression are given for each characteristic to define the characteristic and to harmonize descriptions. Each state of expression is allocated a corresponding numerical note for ease of recording of data and for the production and exchange of the description.

6.2.2 All relevant states of expression are presented in the characteristic.

6.2.3 Further explanation of the presentation of states of expression and notes is provided in document TGP/7 "Development of Test Guidelines".

6.3 *Types of Expression*

An explanation of the types of expression of characteristics (qualitative, quantitative and pseudo-qualitative) is provided in the General Introduction.

6.4 *Example Varieties*

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic.

6.5 Legend

		English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note
1	2	3	4	5	6	7	
		Name of characteristics in English	Nom du caractère en français	Name des Merkmals auf Deutsch	Nombre del carácter en español		
		states of expression	types d'expression	Ausprägungsstufen	tipos de expresión		

- 1 Characteristic number
- 2 (*) Asterisked characteristic – see Chapter 6.1.2
- 3 Type of expression
 QL Qualitative characteristic – see Chapter 6.3
 QN Quantitative characteristic – see Chapter 6.3
 PQ Pseudo-qualitative characteristic – see Chapter 6.3
- 4 Method of observation (and type of plot, if applicable)
 MG, MS, VG, VS – see Chapter 4.1.5
- 5 (+) See Explanations on the Table of Characteristics in Chapter 8.2
- 6 (a)-(c) See Explanations on the Table of Characteristics in Chapter 8.1
- 7 Not applicable

7. Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteres

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/
1. (*)	QN	MS/VG	(+)			
	Plant: height					
	short				Sase LIS02	3
	medium				Momo Sen	5
	tall				Mio Peach Chuchu	7
2.	QN	MS/VG	(+)			
	Plant: number of primary branches					
	few				Shonai Cross Pink	3
	medium				Exe Pink	5
	many				Illumypink	7
3.	PQ	VG	(+)			
	Plant: position of primary branches					
	upper part only				Saga T2go	1
	upper and middle part				Lilac Pink Thumb	2
	throughout				Cherrybee	3
4.	QN	MS/VG				
	Stem: number of nodes					
	few				Cherrybee 3go	3
	medium				Momo Sen	5
	many				Shonai Cross White	7

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/
5.	QN MS/VG	(+) (a)				
	Leaf: length					
	short				Diamond	3
	medium				Momo Sen	5
	long				Sase LIS02	7
6. (*)	QN MS/VG	(+) (a)				
	Leaf: width					
	narrow				Cherrybee 3go	3
	medium				Momo Sen	5
	broad				Komachi White Dress	7
7. (*)	QN MS/VG	(+) (a)				
	Leaf: ratio length/width					
	low				Komachi White Dress	3
	medium				Momo Sen	5
	high				Shonai Cross White	7
8. (*)	QN VG	(a)				
	Leaf: glaucosity					
	absent or weak				Cherrybee	1
	medium				Komachi Green Dress	2
	strong				Momo Sen	3
9. (*)	QN VG	(+) (a)				
	Leaf: intensity of green color					
	light				Saga T2go	1
	medium				Momo Sen	2
	dark				Lilac Pink Thumb	3
10	QN MS/VG	(+)				
	Pedical: length					
	short				Lilac Pink Thumb	3
	medium				Momo Sen	5
	long				Diamond	7

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/
11	QN MS/VG	(+)				
	Calyx: length					
	short				Cherrybee	3
	medium				Momo Sen	5
	long				Asamiyae	7
12	QL VG	(+)				
	Calyx: anthocyanin coloration					
	absent				Light Blue Thumb	1
	present				Cherrybee	9
13	QN MS/VG					
	Flower: number					
	few				Kirara Apricot 2	3
	medium				Momo Sen	5
	many				Mahoroba Peach	7
14 (*)	QL VG	(+)				
	Flower: type					
	single				Momo Sen	1
	double				Piccorosa Pink Picotee	2
15 (*)	QN MS/VG					
	<u>Only varieties with Flower: type: double:</u> Flower: number of petals					
	few				Komachi Green Dress	3
	medium				Diamond	5
	many				Lination Pink Picotee	7
16	PQ VG	(+)				
	Flower: shape					
	circular				Chigusa	1
	pentagon				Azumanoshirabe	2
	star-shaped				Shonai Cross White	3

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/
17 (*)	QN MS/VG	(+)				
	Flower: width					
	narrow				Chigusa	3
	medium				Momo Sen	5
	broad				Rainbow White	7
18	QN MS/VG	(+)				
	Flower: height					
	short				Chigusa	3
	medium				Momo Sen	5
	tall					7
19	QN MS/VG	(+)				
	Flower: ratio of height/width					
	low				Mahoroba Peach	3
	medium				Momo Sen	5
	high				Shonai Cross White	7
20	PQ VG	(+)	(b)			
	Petal: shape					
	elliptic				Shonai Cross Pink	1
	oblanceolate				Bouquet White	2
	obovate				Momo Sen	3
21 (*)	QN MS/VG	(+)	(b)			
	Petal: length					
	short				Komachi White Dress	3
	medium				Momo Sen	5
	long				Suibijin	7
22 (*)	QN MS/VG	(+)	(b)			
	Petal: width					
	narrow				Shonai Cross White	3
	medium				Momo Sen	5
	broad				Suibijin	7

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/
23	(*)	PQ	VG	(+)	(b)			
		Petal: shape of apex						
			acuminate				Lination Pink Picotee	1
			acute					2
			rounded				Momo Sen	3
			flat				Komachi Green Dress	4
			retuse				Piccorosa Pink Picotee	5
24		QN	VG	(+)	(b)			
		Petal : reflexing of margin						
			absent or very weak				Tokyo E1go	1
			weak				Cute Green	2
			medium				Light Blue Thumb	3
			strong				Momo Sen	4
			very strong				Petit Snow	5
25	(*)	QN	VG	(+)	(b)			
		Petal: undulation of margin						
			weak				Momo Sen	3
			medium				Mio Peach Chuchu	5
			strong				Mahoroba Peach	7
26	(*)	QN	VG	(+)	(b)			
		Petal: depth of incisions of margin						
			absent or shallow				Momo Sen	1
			medium				Mio Peach Chuchu	2
			deep					3
			very deep				Sase LIS02	4
27	(*)	PQ	VG		(b), (c)			
		Petal: main color of inner side (exclud part of base)						
		RHS Colour Chart (indicate reference number)						

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/
28	(*)	PQ	VG	(b), (c)			
		Petal: secondary color of <u>inner</u> side					
		RHS Colour Chart (indicate reference number)					
29	(*)	PQ	VG	(+)	(b), (c)		
		Petal: distribution of secondary color of <u>inner</u> side					
		none					1
		at tip				Komachi Kiss	2
		margin				Piccorosa Pink Picotee	3
		medium stripe					4
		distal half				Mahoroba Peach	5
		basal half				Cherrybee 2go	6
		throughout					7
30	(*)	PQ	VG	(+)	(b), (c)		
		Petal: pattern of secondary color of <u>inner</u> side					
		solid				Piccorosa Pink Picotee	1
		flush				Mahoroba Peach	2
		irregular					3
31	(*)	PQ	VG	(+)	(b)		
		Petal: color of base of <u>inner</u> side					
		green				Chigusa	1
		violet				Momo Sen	2
		brown				Sase LIS02	3
32	(*)	PQ	VG	(b), (c)			
		Petal: main color of <u>outer</u> side					
		RHS Colour Chart (indicate reference number)					
33		QL	VG	(+)			
		Style: anthocyanin coloration					
		absent				Momo Sen	1
		present				Cherrybee 2go	9

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/
34	QN	MG/VG	(+)				
	Only seed -propagated varieties: Time of beginning of flowering						
	early					Cherrybee 3go	3
	medium					Mahoroba Yellow	5
	late					Saga T2go	7

8. Explanations on the Table of Characteristics


8.1 *Explanations covering several characteristics*

Characteristics containing the following key in the Table of Characteristics should be examined as indicated below:

- (a) Observations should be made on the upper side of fully developed leaves from the middle third of a stem.
- (b) Observations on double flowers should be made on a petal from the outermost whorl.
- (c) The main color is the color with the largest area excluding the color at base. The secondary color is the color with the second largest area excluding the color at base. In cases where the areas of the main and secondary color are too similar to decide which color has the largest area, the darker color is considered to be the main color.

8.2 *Explanations for individual characteristics*

Ad. 1: Plant: height




Ad. 2: Plant: number of primary branches

See Ad. 1

Ad. 3: Plant: position of primary branches

See Ad. 1

Ad. 5: Leaf: length



Ad. 6: Leaf: width

See Ad. 5

Ad. 7: Leaf: ratio length/width



3
low



5
medium




7
high

Ad. 9: Leaf: intensity of green color

To be observed on the upper side of a leaf after removing the glaucosity.

Ad. 10: Pedicel: length



Ad. 11: Calyx: length



Ad. 12: Calyx: anthocyanin coloration



1
absent



9
present

Ad. 14: Flower: type

Single varieties have only five petals.



1
single



2
double

Ad. 16: Flower: shape



1
circular




2
pentagon



3
star-shaped

Ad. 17: Flower: width



Ad. 18: Flower: height

See Ad. 17

Ad. 19: Flower: ratio of height/width



3
low



5
medium



7
high

Ad. 20: Petal: shape



1
elliptic




2
oblongate



3
obovate

Ad. 21: Petal: length




Ad. 22: Petal: width

See Ad. 21


Ad. 23: Petal: shape of apex




Ad. 24: Petal : reflexing of margin




Ad. 25: Petal: undulation of margin




Ad. 26: Petal: depth of incisions of margin




Ad. 29: Petal: distribution of secondary color of inner side



Ad. 30: Petal: pattern of secondary color of inner side



Ad. 31: Petal: color of base of inner side



Ad. 33: Style: anthocyanin coloration



Ad. 34: Only seed -propagated varieties: Time of beginning of flowering

Observation should be made on the time of the first flower blooming in 50% of plants.

8.3 Unless otherwise indicated, all observations should be made at the time of full flowering.

9. Literature

Kiyoshi Okawa, 1992: Eustoma (Torukogikyo) Seibundo-Shinkosha Co., Tokyo, JP.

10. Technical Questionnaire

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
		Application date: (not to be filled in by the applicant)
TECHNICAL QUESTIONNAIRE to be completed in connection with an application for plant breeders' rights		
1. Subject of the Technical Questionnaire		
1.1	Botanical name	<input type="text" value="Eustoma exaltatum (L.) Salisb. ex G. Don subsp. russellianum (Hook.) Kartesz"/>
1.2	Common name	<input type="text" value="Eustoma, Lisianthus"/>
2. Applicant		
	Name	<input type="text"/>
	Address	<input type="text"/>
	Telephone No.	<input type="text"/>
	Fax No.	<input type="text"/>
	E-mail address	<input type="text"/>
	Breeder (if different from applicant)	<input type="text"/>
3. Proposed denomination and breeder's reference		
	Proposed denomination (if available)	<input type="text"/>
	Breeder's reference	<input type="text"/>

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

#4. Information on the breeding scheme and propagation of the variety

4.1 Breeding scheme

Variety resulting from:

4.1.1 Crossing

(a) controlled cross

(please state parent variety)

(.....) x (.....)

female parent male parent

(b) partially known cross

(please state known parent variety(ies))

(.....) x (.....)

female parent male parent

(c) unknown cross

4.1.2 Mutation

(please state parent variety)

4.1.3 Discovery and development

(please state where and when discovered and how developed)

4.1.4 Other

(Please provide details)

Authorities may allow certain of this information to be provided in a confidential section of the Technical Questionnaire.

4.2 Method of propagating the variety

4.2.1 Seed-propagated varieties

- (a) Self-pollination []
- (b) Cross-pollination []
- (c) Hybrid []
- (d) Other (please provide details) []

4.2.2 Vegetative propagation

- (a) Cuttings []
- (b) *In vitro* propagation []
- (c) Other (state method) []

4.2.3 Other []
(Please provide details)

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the note which best corresponds).

Characteristics	Example Varieties	Note
5.1 Plant: height (1)		
very short		1 []
very short to short		2 []
short	Sase LIS02	3 []
short to medium		4 []
medium	Momo Sen	5 []
medium to tall		6 []
tall	Mio Peach Chuchu	7 []
tall to very tall		8 []
very tall		9 []
5.2 Flower: type (14)		
single	Momo Sen	1 []
double	Piccorosa Pink Picotee	2 []
5.3 Flower: width (17)		
very narrow		1 []
very narrow to narrow		2 []
narrow	Chigusa	3 []
narrow to medium		4 []
medium	Momo Sen	5 []
medium to broad		6 []
broad	Rainbow White	7 []
broad to very broad		8 []
very broad		9 []

Characteristics	Example Varieties	Note
5.4 Petal: main color of <u>inner</u> side (exclud part of base) (27)		
white		1 []
light green		2 []
yellow		3 []
orange		4 []
pink		5 []
red		6 []
purple		7 []
blue purple		8 []
5.5 Petal: secondary color of <u>inner</u> side (28)		
white		1 []
light green		2 []
yellow		3 []
orange		4 []
pink		5 []
red		6 []
purple		7 []
blue purple		8 []
5.6 Petal: distribution of secondary color of <u>inner</u> side (29)		
none		1 []
at tip	Komachi Kiss	2 []
margin	Piccorosa Pink Picotee	3 []
medium stripe		4 []
distal half	Mahoroba Peach	5 []
basal half	Cherrybee 2go	6 []
throughout		7 []
5.7 Petal: color of base of <u>inner</u> side (31)		
green	Chigusa	1 []
violet	Momo Sen	2 []
brown	Sase LIS02	3 []

Characteristics	Example Varieties	Note
5.8 (34)	Only seed -propagated varieties: Time of beginning of flowering	
very early		1 []
very early to early		2 []
early	Cherrybee 3go	3 []
early to medium		4 []
medium	Mahoroba Yellow	5 []
medium to late		6 []
late	Saga T2go	7 []
late to very late		8 []
very late		9 []

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

6. Similar varieties and differences from these varieties

Please use the following table and box for comments to provide information on how your candidate variety differs from the variety (or varieties) which, to the best of your knowledge, is (or are) most similar. This information may help the examination authority to conduct its examination of distinctness in a more efficient way.

Denomination(s) of variety(ies) similar to your candidate variety	Characteristic(s) in which your candidate variety differs from the similar variety(ies)	Describe the expression of the characteristic(s) for the similar variety(ies)	Describe the expression of the characteristic(s) for your candidate variety
<i>Example</i>	<i>Plant: height</i>	<i>short</i>	<i>medium</i>
<p>Comments:</p>			

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

#7. Additional information which may help in the examination of the variety

7.1 In addition to the information provided in sections 5 and 6, are there any additional characteristics which may help to distinguish the variety?

Yes No

(If yes, please provide details)

7.2 Are there any special conditions for growing the variety or conducting the examination?

Yes No

(If yes, please provide details)

7.3 Other information

A representative color photograph of the variety displaying its main distinguishing feature(s), should accompany the Technical Questionnaire. The photograph will provide a visual illustration of the candidate variety which supplements the information provided in the Technical Questionnaire.

The key points to consider when taking a photograph of the candidate variety are:

- Indication of the date and geographic location
- Correct labeling (breeder's reference)
- Good quality printed photograph (minimum 10 cm x 15 cm) and/or sufficient resolution electronic format version (minimum 960 x 1280 pixels)"

Further guidance on providing photographs with the Technical Questionnaire is available in document TGP/7 "Development of Test Guidelines", Guidance Note 35 (<http://www.upov.int/tgp/en/>).

[The link provided may be deleted by members of the Union when developing authorities' own test guidelines.]

Authorities may allow certain of this information to be provided in a confidential section of the Technical Questionnaire.

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

8. Authorization for release

(a) Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health?

Yes [] No []

(b) Has such authorization been obtained?

Yes [] No []

If the answer to (b) is yes, please attach a copy of the authorization.

9. Information on plant material to be examined or submitted for examination

9.1 The expression of a characteristic or several characteristics of a variety may be affected by factors, such as pests and disease, chemical treatment (e.g. growth retardants or pesticides), effects of tissue culture, different rootstocks, scions taken from different growth phases of a tree, etc.

9.2 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If the plant material has undergone such treatment, full details of the treatment must be given. In this respect, please indicate below, to the best of your knowledge, if the plant material to be examined has been subjected to:

(a)	Microorganisms (e.g. virus, bacteria, phytoplasma)	Yes []	No []
(b)	Chemical treatment (e.g. growth retardant, pesticide)	Yes []	No []
(c)	Tissue culture	Yes []	No []
(d)	Other factors	Yes []	No []

Please provide details for where you have indicated "yes".

.....

10. I hereby declare that, to the best of my knowledge, the information provided in this form is correct:

Applicant's name

Signature Date

[End of document]